



6% boron (B), 0.3% copper (Cu) chelated by EDTA, 0.3% iron (Fe) as complex, 12% manganese (Mn), 5.5% zinc (Zn)



## A combination of key nutrients for foliar application to a range of crops

## ADVANTAGES

- Betatrel is a complete trace element fertiliser
- Betatrel has been carefully formulated for the needs of the plant.
- Soluble powder. It does not block the nozzles.
- High boron and manganese content.
- Provides the correct balance of nutrient levels and ratios to satisfy plant demands.
- Phosyn has over 35 years of experience in the preparation and use of high quality trace elements.





Always read the label before using the product. Utilise soil and leaf analysis to check your crop's nutritional status. For more information on Phosyn and Phosyn products visit: www.phosyn.com.

## **APPLICATION RATES**

Beans: 3 kg/ha at the 6 leaf stage and again 10 to 14 days later. Water rate: 200 l/ha.

Cereals: 2.5 kg/ha from the two leaf stage to first node detectable (Zadoks G.S. 12 to31). Water rate: 200 l/ha minimum.

Groundnuts/Peanuts: 5 kg/ha at the 4 to 6 leaf stage and again 10 to 14 days later. Water rate: 200 l/ha.

Maize: 3 to 5 kg/ha at 4 to 8 leaf stage and 10 to 14 days later. Water rate: 200 l/ha.

**Onion:** Two applications of 3 kg/ha sprayed at the 6 leaf stage and again 10 to 14 days later. Water rate: 500 l/ha.

Pivot Application: Arable Crops for which there is an Approved Foliar Spray Recommendation: Apply at the same rates and timings as for foliar application. Water rate: 50,000 to 100,000 l/ha.

Sunflower: 5 kg/ha applied at 2 to 8 pairs of leaves, with a repeat application 2 weeks later. Water rate: 200 l/ha.



Distributor:

**TriaChem (T) Limited** P.O.Box.2750, Arusha-Tanzania Tel :+255 247250 2446/2503485 Fax :+255 27254 8774 web : www.triachem.com