

# CALMABON

## SPECIAL FERTILISER / SPESIALE MISSTOF

### LIQUID / VLOEISTOF

N – 132 g/l  
Ca – 133 g/l  
Mg – 26 g/l  
B – 2289 mg/l  
Mo – 2275 mg/l

WS/CL Free WO/CL Vry

MISSTOF GR 1 FERTILISER  
Reg. No. / Nr. K5272

### GEBRUIKSAANWYSINGS

Dit word algemeen aanvaar dat die verhouding tussen Kalsium en Magnesium in plante en in die grond 'n kation samestelling van ongeveer 65% teenoor 25% op me% basis moet wees. Calmabon voldoen uitmuntend aan hierdie vereiste. 'n Oormaat Kalsium toediening kan 'n Boor tekort veroorsaak. Die hoeveelheid Boor in Calmabon sal hierdie probleem tenwerk.

**BELANGRIK:** Raadpleeg 'n gekwalifiseerde agronoom of landboukundige voordat hierdie misstof toegedien word oor die noodsaaklikheid van die gebruik daarvan.

### TOEDIENINGS

#### Aanwysings:

Blaarvoeding : meng 5 liter met 200 liter water

Lugbespuiting : 5 liter/30 liter water

Mikro- en drupbesproëing:

GEWAS	TOEDIENING		OPMERKING
	Aantal	l/ha	
Appels	4	6	Dien toe net voor blomval.
Groente	3	3	Dien toe net voor en tydens vrugvorming.
Aartappels	3	3	Dien toe net voor knolvorming.
Tamaties	3	3	Dien toe net na eerste blomme gevorm het. (Moenie oormatig toedien nie)
Sonneblomme	1	3	Dien toe net voor blomme gevorm het.

### DIRECTIONS FOR USE

It is generally accepted that the ratio between Calcium and Magnesium in the soil and plant should be a cation composition of 65% to 25% on a me% basis. Calmabon is ideally suited to meet these requirements. An excess of Calcium may result in a Boron deficiency. The Boron in Calmabon will solve this problem.

**IMPORTANT:** Before applying this fertiliser consult a qualified agronomist or agricultural advisor on the necessity of using it.

### APPLICATIONS

#### Instructions:

Foliar feeding : mix 5 litres with 200 litres water

Aerial spray : 5 litres/30 litres water

Micro and Drip irrigation:

CROP	APPLICATION		REMARKS
	No. of	l/ha	
Apples	4	6	Apply just before first flower drop.
Vegetables	3	3	Apply just before and during fruit formation.
Potatoes	3	3	Apply just before tuber formation.
Tomatoes	3	10 litre	Apply after first flower formation. (Do not over apply)
Sunflowers	1	3	Apply with first flower formation.

VERVAARDIG EN VERSPREI DEUR / MANUFACTURED AND DISTRIBUTED BY

# HYGROTECH

PROPERTIES (PTY) LTD

Reg. No. 1984/00638/07

1 Gerhard Braak Street, Pyramid, 0120

PO Box 17220, Pretoria North, 0116

Tel (012) 545 8000 • Fax (012) 545 0150