

# KIC-START

## Vloeistof / Liquid 7:3:1 Oplossing / Solution

Stikstof / Nitrogen (N)	220	g/L
Fosfor / Phosphorus (P)	91	g/L
Kalium / Potassium (K)	30	g/L
Calcium / Kalsium (Ca)	10	g/L
Magnesium / Magnesium (Mg)	2.5	g/L
Yster / Iron (Fe)	1005	mg/L
Koper / Copper (Cu)	510	mg/L
Sink / Zinc (Zn)	510	mg/L
Mangaan / Manganese (Mn)	510	mg/L
Boor / Boron (B)	261	mg/L
Molibdeen / Molybdenum (Mo)	8.4	mg/L
Ouksiene / Auxins	5.0	mg/L

FERTILIZER GROUP 1  
Reg. Nr. K5442

MISSTOF GROEP 1  
Reg. No. K5442

Vervaardig en versprei deur / Manufactured and distributed by:

## HYGROTECH

SEED (PTY) LTD / SAAD (EDMS) BPK

Reg. No. 84/00638/07

1 Gerhard Braak St, Pyramid, 0120

P.O. Box 17220, Pretoria North, 0116

Tel (012) 545 8 000 · Fax (012) 545 0150

### DIRECTIONS FOR USE

KIC-START is a water soluble nutrient solution with chelated cation micro elements which can be used as a root drench when transplanting seedlings or when establishing trees and ornamental shrubs.

KIC-START is compatible with most fungicides and pesticides except those which contain large amounts of copper, sulphur and alkaline materials such as lime sulphur or Bordeaux mixtures.

KIC-START is formulated to stimulate growth during first crucial weeks after transplanting.

To enhance the compatibility of multiple tank mixes SPRAY-AIDE\* should be used. If KIC-START is used in waters with an abnormally high or low pH, a buffer such as HYGROBUFF 4\* should be used.

SPRAY-AIDE\*                      L3308 ACT 36/1947  
HYGROBUFF 4\*                    L5512 ACT 36/1947

### KIC-START SOLUTION

The recommended dosage of KIC-START is 2 litre of product per 400 litre of water (i.e. 5ml/litre water or 500ml/ 100 litre water).

KIC-START solution can be applied into the planting hole just before transplanting at a rate of 250ml per seedling.

KIC-START solution can be used as a soil drench directly after transplant at a rate of 250ml per seedling.

KIC-START solution can be applied to the planting hole of ornamental and forestry plants at a rate of 500-750 ml / plant.

KIC-START can also be applied through the drip irrigation system: Apply the concentrate at the end of an irrigation cycle within the first few days after transplanting. Apply at a rate of 2L per hectare.

### GEBRUIKSAANWYSINGS

KIC-START is 'n wateroplosbare plantvoeding met gecheleerde kation mikro elemente wat as 'n wortelbehandeling toegedien word by die uitplant van saailinge of by die vestiging van bome en ornamentele struik.

KIC-START is verenigbaar met meeste swam- en insekdoders, behalwe dié wat groot hoeveelhede koper, swawel of alkaliese verbindings bevat soos byvoorbeeld kalkswawel of Bordeaux mengsels.

KIC-START word gebruik om groei te stimuleer gedurende die eerste paar weke na uitplanting.

Om die mengbaarheid in die tenk te verbeter moet SPRAY-AIDE\* bygevoeg word. HYGROBUFF 4\* kan ook bygevoeg word indien water met 'n hoë of lae pH gebruik word.

SPRAY-AIDE\*                      L3308 ACT 36/1947  
HYGROBUFF 4\*                    L5512 ACT 36/1947

### KIC-START OPLOSSING

Die aanbevele toedieningshoeveelheid vir KIC-START is 2 liter konsentraat per 400 liter water (b.v. 5ml/liter water of 500ml/100 liter water).

KIC-START oplossing word teen 'n dosis van 250ml per saailing direk in die gat toegedien tydens uitplanting.

KIC-START kan direk na uitplanting as 'n wortelbehandeling toegedien word en kan ook rondom elke saailing gegooi word teen 'n dosis van 250ml per saailing.

KIC-START kan teen 'n dosis van 500-750 ml per plantgat aan ornamentele struik en bosbou bome tydens uitplanting toegedien word.

KIC-START kan ook deur die drupper toegedien word. Dien die konsentraat toe aan die einde van die besproeiings siklus binne die eerste paar dae nadat uitgeplant is. Dien dit toe teen 'n dosis van 2L per hektaar.