


Brigade

025 EC Insecticide/Miticide



Member of Pesticides Chemical Association of Kenya Membership No. PCA.128

Bayer East Africa Agrochemicals 

Directions for use

Measure the required quantity of concentrate for the spray tank volume, add it to a small quantity of water. In the sprayer, stir the mixture and fill sprayer tank to desired volume while stirring. Ensure full cover of all target surface when spraying.

CROP	PEST	USE DIRECTIONS
Citrus	Citrus scales, Aphids, Mealybug, Psyllid, Citrus bud and rust mites, leafminer and Citrus black fly	Apply 25ml product in 10 litre sprayer Preharvest period 72 hours
Vegetables (including: Cabbages, Kales, Beans, Peas, Peppers, Brinjals, Okra, Tomato, Onions Melons etc.	Heliothis Bollworm, Diamond back moth, Aphids, Thrips, Cabbage Sawfly, Whitefly, Maruca Pod borer, Striped blister beetle Red Spider mite (Tetranychus spp) Whitefly	Apply 16ml product in 10 litre sprayer Apply 25ml product per 10 Litre sprayer Preharvest period 72 hours
Ornamentals	Heliothis bollworm, Aphids Soft scale, Cushiony scale and Mealybug Spider mites	Apply 16ml per 10 litre sprayer Apply 40 ml/10 litre sprayer

CROP	PEST	USE DIRECTIONS
		Warning: Always check for varietal tolerance before large scale application.
General soil pests	<i>Agrotis</i> (cut worms)	Spray seedbed evenly with 16ml product per 10 litre sprayer. The treatment also effectively controls soil beetles.
Coffee	Cottony scale (<i>Icerya pattersoni</i>)	Apply 12ml in 10 litre sprayer Preharvest period 48 hours.
Wheat Barley	Aphids, vector of BYDV	Apply 12ml in 10 litre sprayer. Preharvest period 14 days
Cotton	Cotton pest complex including <i>Heliothis</i> spp. <i>Lygus</i> spp. Pink bollworm, Aphids, Leafworm etc.	Apply 25ml in 10 litres sprayer
	Spider mites	Apply 40ml in 10 litre sprayer
Maize	Maize stalk borer	Apply 12ml in 10 litre sprayer

First Aid:

Eye Splashes: Brigade is a moderate irritant. If splashed onto the eyes flush out with clean water for at least 15 minutes and then obtain immediate medical attention.

Splashes on skin:

Remove contaminated clothing and wash skin immediately with plenty of water and soap.

If swallowed:

Do not induce victim to vomit as aromatic solvents may damage the lungs. Obtain immediate medical attention and carry the container/label to the doctor.

Toxicology:**Guide to Doctor:**

Brigade is synthetic Pyrethroid containing aromatic solvents. If swallowed perform gastric lavage taking care to prevent aspiration of gastric contents. Therefore treat symptomatically. No antidote. **Symptoms of Poisoning:** Weakness and subdued behaviour, sweating, blurred vision, dehydration, restlessness, difficult breathing.

Notice to users:

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

Warranty Clause:

All chemicals manufactured by us are tested, and if properly applied, are suitable for the purpose recommended

on the label. However, FMC Corporation cannot exercise full control over their application by the user, we are therefore unable to accept any responsibility for damage arising either from their direct or indirect application for use.

Brigade is the registered trade mark of FMC Corporation U.S.A.

Brigade is manufactured and formulated by FMC Corporation and distributed in Kenya by Bayer East Africa Limited, Lungu Lungu Road, P.O. Box 30321, Telephone 531272-5, Nairobi.